

HELMHOLTZ
MUNICH

AIH Institute of AI for Health

Topological Machine Learning: The (W)Hole Truth

Lecture 1

Bastian Rieck (@Pseudomanifold)

Preliminaries




Do you have feedback or any questions? Write to bastian.riek@helmholtz-muenchen.de or reach out to [@Pseudomanifold](https://twitter.com/Pseudomanifold) on Twitter. You can find the slides and additional information with links to more literature here:

<https://heidelberg.topology.rocks>




Philosophy

Computational topology is all about finding *expressive* and *computable* invariants to distinguish between different spaces.




What is computational topology?






What is computational topology?



What is computational topology?



What is computational topology?




Which qualities of the sphere make it *different* from the torus?

Betti numbers

The d^{th} Betti number counts the number of d -dimensional holes. It can be used to distinguish between spaces.

β_0 Connected components
 β_1 Tunnels
 β_2 Voids

Space	β_0	β_1	β_2
Point	1	0	0
Cube	1	0	1
Sphere	1	0	1
Torus	1	2	1



Simplicial complexes

Definition

A set family K is called an *abstract simplicial complex* if, for every set $\sigma \in K$ and each non-empty subset $\tau \subseteq \sigma$, we also have $\tau \in K$.

Terminology

The elements of a simplicial complex K are called *simplices*. Notice that a k -simplex consists of $k + 1$ vertices.


Visualisation

We will typically draw simplicial complexes like graphs, with the understanding that the positions of vertices are *arbitrary* unless some geometric component is involved.

Simplicial complexes

Example


Simplicial complexes can be decomposed into their skeletons, which only contain simplices of a certain dimension.



Simplicial complexes

Example


Simplicial complexes can be decomposed into their skeletons, which only contain simplices of a certain dimension.



Simplicial complexes

Example


Simplicial complexes can be decomposed into their skeletons, which only contain simplices of a certain dimension.



Simplicial complexes

Example


Simplicial complexes can be decomposed into their skeletons, which only contain simplices of a certain dimension.



Simplicial complexes

Non-example

This is *not* a simplicial complex because some higher-dimensional simplices do not intersect in a lower-dimensional one!



Simplicial complexes

More examples

☆ Graphs can be considered (low-dimensional) simplicial complexes.

Simplicial complexes

More examples

- ☆ Graphs can be considered (low-dimensional) simplicial complexes.
- ☆ Simplicial complexes can be obtained from point clouds (more about this later).

Simplicial complexes

More examples

- ☆ Graphs can be considered (low-dimensional) simplicial complexes.
- ☆ Simplicial complexes can be obtained from point clouds (more about this later).
- ☆ *Hypergraphs* can be converted to simplicial complexes.

To calculate something *on* simplicial complexes,
we need some group theory.

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

- 1 The operation is *closed*, i.e. $a \cdot b \in G$ for $a, b \in G$.

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

- 1 The operation is *closed*, i.e. $a \cdot b \in G$ for $a, b \in G$.
- 2 The operation is *associative*, i.e. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for $a, b, c \in G$.

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

- 1 The operation is *closed*, i.e. $a \cdot b \in G$ for $a, b \in G$.
- 2 The operation is *associative*, i.e. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for $a, b, c \in G$.
- 3 There is an *identity element* $e \in G$ such that $e \cdot a = a \cdot e = a$ for $a \in G$.

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

- 1 The operation is *closed*, i.e. $a \cdot b \in G$ for $a, b \in G$.
- 2 The operation is *associative*, i.e. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for $a, b, c \in G$.
- 3 There is an *identity element* $e \in G$ such that $e \cdot a = a \cdot e = a$ for $a \in G$.
- 4 Each $a \in G$ has an *inverse element* $a^{-1} \in G$ such that $a \cdot a^{-1} = e = a^{-1} \cdot a$.

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

- 1 The operation is *closed*, i.e. $a \cdot b \in G$ for $a, b \in G$.
- 2 The operation is *associative*, i.e. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for $a, b, c \in G$.
- 3 There is an *identity element* $e \in G$ such that $e \cdot a = a \cdot e = a$ for $a \in G$.
- 4 Each $a \in G$ has an *inverse element* $a^{-1} \in G$ such that $a \cdot a^{-1} = e = a^{-1} \cdot a$.

Groups

Definition

A *group* is a set G with a binary operation \cdot that combines two elements to yield another one, such that (G, \cdot) has the following properties:

- 1 The operation is *closed*, i.e. $a \cdot b \in G$ for $a, b \in G$.
- 2 The operation is *associative*, i.e. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for $a, b, c \in G$.
- 3 There is an *identity element* $e \in G$ such that $e \cdot a = a \cdot e = a$ for $a \in G$.
- 4 Each $a \in G$ has an *inverse element* $a^{-1} \in G$ such that $a \cdot a^{-1} = e = a^{-1} \cdot a$.

The operation \cdot is not required to be commutative. In general, $a \cdot b = b \cdot a$ is *not* required to hold. However, the groups that we will encounter are commutative!

Groups

Examples and non-examples

☆ The set with only two elements and addition modulo 2 is group, called \mathbb{Z}_2 or \mathbb{F}_2 .¹

¹It is actually also a *field*, the smallest non-trivial field.

Groups

Examples and non-examples

- ☆ The set with only two elements and addition modulo 2 is group, called \mathbb{Z}_2 or \mathbb{F}_2 .¹
- ☆ The set of integers \mathbb{Z} with the usual addition is a group, denoted by $(\mathbb{Z}, +)$.

¹It is actually also a *field*, the smallest non-trivial field.

Groups

Examples and non-examples

- ☆ The set with only two elements and addition modulo 2 is group, called \mathbb{Z}_2 or \mathbb{F}_2 .¹
- ☆ The set of integers \mathbb{Z} with the usual addition is a group, denoted by $(\mathbb{Z}, +)$.
- ☆ The set of \mathbb{R} -valued quadratic matrices with elementwise addition is a group.

¹It is actually also a *field*, the smallest non-trivial field.

Groups

Examples and non-examples

- ☆ The set with only two elements and addition modulo 2 is group, called \mathbb{Z}_2 or \mathbb{F}_2 .¹
- ☆ The set of integers \mathbb{Z} with the usual addition is a group, denoted by $(\mathbb{Z}, +)$.
- ☆ The set of \mathbb{R} -valued quadratic matrices with elementwise addition is a group.
- ☆ The set of \mathbb{R} -valued quadratic matrices with non-zero determinant together with matrix multiplication is a group.

¹It is actually also a *field*, the smallest non-trivial field.

Groups

Examples and non-examples

- ☆ The set with only two elements and addition modulo 2 is group, called \mathbb{Z}_2 or \mathbb{F}_2 .¹
- ☆ The set of integers \mathbb{Z} with the usual addition is a group, denoted by $(\mathbb{Z}, +)$.
- ☆ The set of \mathbb{R} -valued quadratic matrices with elementwise addition is a group.
- ☆ The set of \mathbb{R} -valued quadratic matrices with non-zero determinant together with matrix multiplication is a group.
- ☆ The natural numbers \mathbb{N} with addition do *not* form a group (why?).

¹It is actually also a *field*, the smallest non-trivial field.

Back to simplicial complexes...

Chain groups

Free abelian groups

Definition

Given a simplicial complex K , the p^{th} chain group C_p of K consists of all combinations of p -simplices in the complex. Coefficients are in \mathbb{Z}_2 , hence all elements of C_p are of the form $\sum_j \sigma_j$, for $\sigma_j \in K$. The group operation is addition with \mathbb{Z}_2 coefficients.

Chain groups

Free abelian groups

Definition

Given a simplicial complex K , the p^{th} chain group C_p of K consists of all combinations of p -simplices in the complex. Coefficients are in \mathbb{Z}_2 , hence all elements of C_p are of the form $\sum_j \sigma_j$, for $\sigma_j \in K$. The group operation is addition with \mathbb{Z}_2 coefficients.

\mathbb{Z}_2 is convenient for implementation reasons because *addition* can be implemented as *symmetric difference*. Other choices are possible!

Chain groups

Free abelian groups


Definition

Given a simplicial complex K , the p^{th} chain group C_p of K consists of all combinations of p -simplices in the complex. Coefficients are in \mathbb{Z}_2 , hence all elements of C_p are of the form $\sum_j \sigma_j$, for $\sigma_j \in K$. The group operation is addition with \mathbb{Z}_2 coefficients.

\mathbb{Z}_2 is convenient for implementation reasons because *addition* can be implemented as *symmetric difference*. Other choices are possible!

We need chain groups to algebraically express the concept of a *boundary*.


Simplicial chains



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

$$\star \{a, b\}$$

Simplicial chains




Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

☆ $\{a, b\}$

☆ $\{a, c\}$


Simplicial chains



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

- ☆ $\{a, b\}$
- ☆ $\{a, c\}$
- ☆ $\{b, c\}$


Simplicial chains



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

- ☆ $\{a, b\}$
- ☆ $\{a, c\}$
- ☆ $\{b, c\}$
- ☆ $\{a, b\} + \{a, c\}$


Simplicial chains



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

- ☆ $\{a, b\}$
- ☆ $\{a, c\}$
- ☆ $\{b, c\}$
- ☆ $\{a, b\} + \{a, c\}$
- ☆ $\{a, b\} + \{b, c\}$


Simplicial chains



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

- ☆ $\{a, b\}$
- ☆ $\{a, c\}$
- ☆ $\{b, c\}$
- ☆ $\{a, b\} + \{a, c\}$
- ☆ $\{a, b\} + \{b, c\}$
- ☆ $\{a, c\} + \{b, c\}$

Simplicial chains



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. Some valid simplicial 1-chains of K are:

- ☆ $\{a, b\}$
- ☆ $\{a, c\}$
- ☆ $\{b, c\}$
- ☆ $\{a, b\} + \{a, c\}$
- ☆ $\{a, b\} + \{b, c\}$
- ☆ $\{a, c\} + \{b, c\}$
- ☆ $\{b, c\} + \{a, c\} + \{a, b\}$

Group homomorphism

Definition

A function $f: A \rightarrow B$ between groups (A, \cdot) and $(B, *)$ is called *homomorphism* if $f(x \cdot y) = f(x) * f(y)$ for all $x, y \in A$.

Boundary homomorphism

Definition

Given a simplicial complex K , the p^{th} boundary homomorphism is a function that assigns each simplex $\sigma = \{v_0, \dots, v_p\} \in K$ to its *boundary*:

$$\partial_p \sigma = \sum_i \{v_0, \dots, \widehat{v}_i, \dots, v_p\}$$


In the equation above, \widehat{v}_i indicates that the set does *not* contain the i^{th} vertex. The function $\partial_p: C_p \rightarrow C_{p-1}$ is thus a homomorphism between the chain groups.

Caveat

With other coefficients, the boundary homomorphism is slightly more complex, involving alternating signs for the different terms. Over \mathbb{Z}_2 , signs can be ignored.

Boundary homomorphism

Example



Let $K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$. The boundary of the triangle is non-trivial:

$$\partial_2 \{a, b, c\} = \{b, c\} + \{a, c\} + \{a, b\}$$

The boundary of its edges is trivial, though, because duplicate simplices cancel each other out:

$$\begin{aligned} \partial_1 (\{b, c\} + \{a, c\} + \{a, b\}) &= \{c\} + \{b\} + \{c\} + \{a\} + \{b\} + \{a\} \\ &= 0 \end{aligned}$$

Chain complex

For all p , we have $\partial_{p-1} \circ \partial_p = 0$: *Boundaries do not have a boundary themselves*. This leads to the *chain complex*:

$$0 \xrightarrow{\partial_{n+1}} C_n \xrightarrow{\partial_n} C_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \xrightarrow{\partial_0} 0$$

Kernel and image

Definition

The *kernel* of a homomorphism $f: A \rightarrow B$ is the set of all elements that are mapped to the zero element, i.e. $\ker f := \{a \in A \mid f(a) = 0\} \subseteq A$. The *image* of f is the set of all its outputs, i.e. $\operatorname{im} f := \{f(a) \mid a \in A\} \subseteq B$.

Cycle and boundary groups

Cycle group $Z_p = \ker \partial_p$

Boundary group $B_p = \text{im } \partial_{p+1}$

We have $B_p \subseteq Z_p$ in the group-theoretical sense. In other words, every boundary is also a cycle.

(The fact that these sets are groups is a consequence of some deep theorems in group theory!
Unfortunately, we cannot cover all of these things here...)

Normal subgroup and quotient group

Normal subgroup

Let G be a group and N be a subgroup. N is a *normal subgroup* if $gng^{-1} \in N$ for all $g \in G$ and $n \in N$.

For an abelian group, every subgroup is normal!

Definition

Let G be a group and N be a normal subgroup of G . Then the *quotient group* is defined as $G/N := \{gN \mid g \in G\}$, partitioning G into equivalence classes.

Quotient groups

Example

$2\mathbb{Z} \subseteq \mathbb{Z}$ is the subgroup of \mathbb{Z} defined by being a multiple of 2. Hence, $\mathbb{Z}_2 := \mathbb{Z}/2\mathbb{Z}$ consists of only 0 and 1.

Why quotient groups?

Quotient groups 'reduce' a group by partitioning it into equivalence classes that are defined by another subgroup.

Homology groups & Betti numbers

The p^{th} homology group H_p is a quotient group, defined by ‘removing’ cycles that are boundaries from a higher dimension:

$$H_p = Z_p / B_p = \ker \partial_p / \text{im } \partial_{p+1},$$

With this definition, we may finally calculate the p^{th} Betti number:

$$\beta_p = \text{rank } H_p$$


The rank is a generating set of the smallest cardinality. We will see how to calculate this easily!

Intuition

Calculate all boundaries, remove the boundaries that come from higher-dimensional objects, and count what is left.

Example

Simplicial complex



$$K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}\}$$

Notice that K does not contain the 2-simplex $\{a, b, c\}$. Next, we will see how to calculate the boundary matrix of K and its homology groups!

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.

$$M = \begin{matrix} & \begin{matrix} a & b & c & ab & bc & ac \end{matrix} \\ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix} \end{matrix}$$

$a \bullet$

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.


$a \bullet$ $\bullet b$

$$M = \begin{matrix} & \begin{matrix} a & b & c & ab & bc & ac \end{matrix} \\ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix} \end{matrix}$$

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.




$$M = \begin{pmatrix} a & b & c & ab & bc & ac \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix}$$

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.




$$M = \begin{matrix} & \begin{matrix} a & b & c & ab & bc & ac \end{matrix} \\ \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix} \end{matrix}$$

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.




$$M = \begin{pmatrix} a & b & c & ab & bc & ac \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix}$$

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.




$$M = \begin{pmatrix} a & b & c & ab & bc & ac \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix}$$

Example

Boundary matrix calculation

Using simplices as a basis of the chain groups, we can represent all boundary relations in a single matrix, the *boundary matrix*.



$$M = \begin{matrix} & \begin{matrix} a & b & c & ab & bc & ac \end{matrix} \\ \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \begin{matrix} a \\ b \\ c \\ ab \\ bc \\ ac \end{matrix} \end{matrix}$$

Example

Dimension 0

To compute H_0 , we need to calculate $Z_0 = \ker \partial_0$ and $B_0 = \text{im } \partial_1$.

Calculating Z_0

We have $Z_0 = \ker \partial_0 = \text{span}(\{a\}, \{b\}, \{c\})$, because each one of these simplices is mapped to zero. Since we cannot express any one of these simplices as a linear combination of the others, we have $Z_0 = (\mathbb{Z}/2\mathbb{Z})^3$.

Calculating B_0

We have $B_0 = \text{im } \partial_1 = \text{span}(\{a\} + \{b\}, \{b\} + \{c\}, \{a\} + \{c\})$. However, since $\{a\} + \{b\} + \{b\} + \{c\} = \{a\} + \{c\}$, there are only two independent elements, i.e. $\text{im } \partial_1 = \text{span}(\{a\} + \{b\}, \{b\} + \{c\})$. Hence, $B_0 = (\mathbb{Z}/2\mathbb{Z})^2$.

Example

Dimension 0, continued

- ☆ By definition, $H_0 = Z_0/B_0 = (\mathbb{Z}/2\mathbb{Z})^3 / (\mathbb{Z}/2\mathbb{Z})^2 = \mathbb{Z}/2\mathbb{Z}$.
- ☆ Hence, $\beta_0 = \text{rank } H_0 = 1$.

Intuition

Our calculation tells us that the simplicial complex has a *single* connected component!

Example

Dimension 1

To compute H_1 , we need to calculate $Z_1 = \ker \partial_1$ and $B_1 = \text{im } \partial_2$.

Calculating Z_1

We have $Z_1 = \ker \partial_1 = \text{span}(\{a, b\} + \{b, c\} + \{a, c\})$. This is the *only* cycle in K ; we can verify this by inspection or pure combinatorics. Hence, $Z_1 = \mathbb{Z}/2\mathbb{Z}$.

Calculating B_1

There are *no* 2-simplices in K , so $B_1 = \text{im } \partial_2 = \{0\}$.

Example

Dimension 1, continued

☆ By definition, $H_1 = Z_1/B_1 = (\mathbb{Z}/2\mathbb{Z})/\{0\} = \mathbb{Z}/2\mathbb{Z}$.

☆ Hence, $\beta_1 = \text{rank } H_1 = 1$.

Intuition

Our calculation tells us that the simplicial complex has a *single* cycle!



This is one of the few situations in which a ‘division by zero’ is well-defined! By the definition of the quotient group, this means we are *not* removing any elements from the group.

Homology calculations in practice

Smith normal form

Let M be an $n \times m$ matrix with at least one non-zero entry over some field \mathbb{F} . There are invertible matrices S and T such that the matrix product SMT has the form

$$SMT = \begin{pmatrix} b_0 & 0 & 0 & \cdots & 0 \\ 0 & b_1 & 0 & \cdots & 0 \\ 0 & 0 & \ddots & & 0 \\ \vdots & & & b_k & \vdots \\ & & & 0 & \\ & & & & \ddots \\ 0 & \cdots & & & 0 \end{pmatrix},$$

where all the entries b_i satisfy $b_i \geq 1$ and divide each other, i.e. $b_i \mid b_{i+1}$. All b_i are unique up to multiplication by a unit.

Homology calculations in practice

- 1 Calculate boundary operator matrices.

Homology calculations in practice

- 1 Calculate boundary operator matrices.
- 2 Bring each matrix into Smith normal form (similar to Gaussian elimination).

Homology calculations in practice

- 1 Calculate boundary operator matrices.
- 2 Bring each matrix into Smith normal form (similar to Gaussian elimination).
- 3 Read off description of p^{th} homology group.

Homology calculations in practice

- 1 Calculate boundary operator matrices.
- 2 Bring each matrix into Smith normal form (similar to Gaussian elimination).
- 3 Read off description of p^{th} homology group.

Homology calculations in practice

- 1 Calculate boundary operator matrices.
- 2 Bring each matrix into Smith normal form (similar to Gaussian elimination).
- 3 Read off description of p^{th} homology group.

We have:

☆ $\text{rank } Z_p$ is the number of zero columns of the boundary matrix of ∂_p .

Homology calculations in practice

- 1 Calculate boundary operator matrices.
- 2 Bring each matrix into Smith normal form (similar to Gaussian elimination).
- 3 Read off description of p^{th} homology group.

We have:

- ☆ $\text{rank } Z_p$ is the number of zero columns of the boundary matrix of ∂_p .
- ☆ $\text{rank } B_p$ is the number of non-zero rows of the boundary matrix of ∂_{p+1} .

Summary

☆ Homology groups characterise topological objects.

Summary

- ☆ Homology groups characterise topological objects.
- ☆ They can be easily expressed as linear operators.

Summary

- ☆ Homology groups characterise topological objects.
- ☆ They can be easily expressed as linear operators.
- ☆ The calculation of homology groups boils down to linear algebra.